



## 저작자표시 2.0 대한민국

이용자는 아래의 조건을 따르는 경우에 한하여 자유롭게

- 이 저작물을 복제, 배포, 전송, 전시, 공연 및 방송할 수 있습니다.
- 이차적 저작물을 작성할 수 있습니다.
- 이 저작물을 영리 목적으로 이용할 수 있습니다.

다음과 같은 조건을 따라야 합니다:



저작자표시. 귀하는 원저작자를 표시하여야 합니다.

- 귀하는, 이 저작물의 재이용이나 배포의 경우, 이 저작물에 적용된 이용허락조건을 명확하게 나타내어야 합니다.
- 저작권자로부터 별도의 허가를 받으면 이러한 조건들은 적용되지 않습니다.

저작권법에 따른 이용자의 권리는 위의 내용에 의하여 영향을 받지 않습니다.

이것은 [이용허락규약\(Legal Code\)](#)을 이해하기 쉽게 요약한 것입니다.

[Disclaimer](#) 

의학석사 학위논문

Altered Resting State Functional  
Connectivity of Anterior Cingulate  
Cortex and Executive Dysfunction in  
Obsessive-Compulsive Disorder

강박증에서 앞쪽 띠피질을 중심으로  
한 휴지기 기능적 뇌연결성의 변화와  
집행기능 저하

2012년 8월

서울대학교 대학원  
의학과 정신과학 전공  
윤 제 연

A thesis of the Master's degree

강박증에서 앞쪽 띰피질을 중심으로  
한 휴지기 기능적 뇌연결성의 변화와  
집행기능 저하

**Altered Resting State Functional  
Connectivity of Anterior Cingulate  
Cortex and Executive Dysfunction in  
Obsessive-Compulsive Disorder**

August 2012

**The Department of Psychiatry  
Seoul National University  
College of Medicine  
Je-Yeon Yun**

# **Altered Resting State Functional Connectivity of Anterior Cingulate Cortex and Executive Dysfunction in Obsessive-Compulsive Disorder**

**By**

**Je-Yeon Yun, M.D.**

**A thesis submitted in partial fulfillment of the requirement  
for the Degree of Master of Philosophy in medicine  
(Department of Psychiatry) at Seoul National University  
College of Medicine**

**Director: Prof. Jun Soo Kwon, M.D., Ph.D.**

**June 2012**

**Approved by Thesis Committee:**

**Professor \_\_\_\_\_ Chairman**

**Professor \_\_\_\_\_ Vice chairman**

**Professor \_\_\_\_\_**

# 강박증에서 앞쪽 띠피질을 중심으로 한 휴지기 기능적 뇌연결성의 변화와 집행기능 저하

지도교수 권 준 수

이 논문을 의학석사 학위논문으로 제출함

2012 년 04 월

서울대학교 대학원

의학과 정신과학 전공

윤 제 연

윤제연의 의학석사 학위논문을 인준함

2012 년 06 월

위 원 장 (인)

부위원장 (인)

위 원 (인)

# ABSTRACT

**Introduction:** The anterior cingulate cortex (ACC) is one of the prefrontal targets of the cortico-striato-thalamo-cortical (CSTC) loops in obsessive-compulsive disorder (OCD) and has been regarded as a neural correlate of executive dysfunction. We investigated the resting state functional connectivity networks (rs-FCNs) of ACC functional subdivisions with the aim of clarifying the relationship between rs-FCN derangements and executive dysfunction in patients with OCD

**Methods:** Twenty-one patients with OCD were recruited from Seoul National University Hospital and 22 healthy volunteers (HC) underwent a resting state functional magnetic resonance imaging (fMRI) scan. The rs-FCN maps were depicted using five region-of-interest (ROI) seed regions systemically placed throughout the ACC. Correlations between the five neuropsychological test performances and the regional ACC seed-based rs-FCNs with significantly altered functional connectivity strength were examined in between-group analyses.

**Results:** The rs-FCNs showed rostral/caudal functional distinction in both groups. Between-group analyses revealed significantly weaker functional connectivity of the OCD group in the claustrum, thalamus, and inferior parietal lobule (IPL). Stronger rs-FC between ACC Seed S5 and the right claustrum correlated with higher Rey–Osterrieth Complex Figure Test organization scores. The prolonged Trail Making Test part-A reaction time demonstrated a correlation with weaker rs-FC strength between ACC Seed S7 and the right IPL.

**Conclusion:** Altered strengths in the functional connectivity between ACC seeds versus specific brain areas of the CSTC loop may be the neural

correlates of executive dysfunction in OCD. Additionally, the inferior parietal lobule could be one of the important brain regions included in the OCD pathophysiologic loops.

**Keywords:** Anterior cingulate cortex, resting state functional connectivity, functional magnetic resonance imaging, obsessive-compulsive disorder, executive dysfunction

**Student Number:** 2010-23694

# CONTENTS

Abstract-----	i
Contents-----	iii
List of tables and figures-----	iv
List of Abbreviations-----	vii
Introduction-----	1
Materials and Methods-----	6
Results-----	13
Discussion-----	43
References-----	48
Abstract in Korean-----	60



## LIST OF TABLES AND FIGURES

Table 1 Demographic and clinical characteristics of obsessive-compulsive disorder group and healthy control-----	14
Table 2 Neuropsychological test scores of the subjects-----	15
Table 3 Regional difference in resting state functional connectivity network strength between patients with obsessive-compulsive disorder and healthy controls-----	26
Table 4 Significance of diagnosis in predicting strength of positive resting state functional connectivity-----	27
Table 5 Regional difference in negative correlations of resting state functional connectivity network between the patients with obsessive-compulsive disorder and healthy controls-----	30
Table 6 Significance of diagnosis in predicting strength of negative resting state functional connectivity-----	31
Table 7 Correlation between altered regional resting state functional connectivity strength versus Y-BOCS scores in Obsessive-Compulsive disorder subjects-----	33
Table 8 Bivariate correlations between regional positive functional connectivity strength and neuropsychological test performance in Obsessive-Compulsive disorder patients-----	37
Table 9 Bivariate correlations between regional positive functional connectivity strength and neuropsychological test performance in healthy controls-----	38
Table 10 Bivariate correlations between regional negative functional connectivity strength and neuropsychological test performance in Obsessive-Compulsive disorder patients-----	39

Table 11 Bivariate correlations between regional negative functional connectivity strength and neuropsychological test performance in healthy controls-----	40
Figure 1 Positive and negative resting state functional connectivity networks between the S1 ACC seed region and other brain area in Obsessive-Compulsive Disorder group-----	16
Figure 2 Positive and negative resting state functional connectivity networks between the S3 ACC seed region and other brain area in Obsessive-Compulsive Disorder group-----	17
Figure 3 Positive and negative resting state functional connectivity networks between the S5 ACC seed region and other brain area in Obsessive-Compulsive Disorder group-----	18
Figure 4 Positive and negative resting state functional connectivity networks between the S7 ACC seed region and other brain area in Obsessive-Compulsive Disorder group-----	19
Figure 5 Positive and negative resting state functional connectivity networks between the I9 ACC seed region and other brain area in Obsessive-Compulsive Disorder group-----	20
Figure 6 . Positive resting state functional connectivity networks of the 5 ACC seed ROIs in OCD and healthy control. Significant within-group functional connectivity maps of the 5 ACC seeds are placed on 3-dimensional anatomical images-----	23
Figure 7 Significant between-group differences in positive resting state functional connectivity networks of the 5 ACC seed ROIs----	25
Figure 8 Correlation between positive regional functional connectivity strength and Y-BOCS scores in obsessive-compulsive disorder---	34

Figure 9 Correlation between negative regional functional connectivity strength and Y-BOCS scores in obsessive-compulsive disorder----- 35

Figure 10 Correlation between regional functional connectivity strength and neuropsychological test performance in obsessive-compulsive disorder----- 41

## LIST OF ABBREVIATIONS

ACC: anterior cingulate cortex

BA: Brodmann area

BAI: Beck Anxiety Inventory

BDI: Beck Depression Inventory

COWAT: Controlled Oral Word Association Test

CSTC: cortico-striato-thalamo-cortical

DLPFC: dorsolateral prefrontal cortex

DMN: default mode network

DSM-IV: Diagnostic and Statistical Manual of Mental Disorders, 4<sup>th</sup> edition

ERN: error-related negativity

fMRI: functional magnetic resonance imaging

IPL: inferior parietal lobule

IQ: intelligence quotient

K-WAIS: Korean-Wechsler Adult Intelligence Scale

MNI: Montreal Neurologic Institute

OCD: Obsessive-compulsive disorder

OFC: orbitofrontal cortex

RCFT: Rey-Osterrieth Complex Figure Test

RS-FCN: resting state functional connectivity network

ROI: Region of Interest

PCC: posterior cingulated cortex

SCID-IV: Structured Clinical Interview for DSM-IV

SCID-NP: Structured Clinical Interview for DSM-IV Non-patient Edition

SCWT: Stroop Color-Word Test

SPM: Statistical Parametric Mapping

TMT: Trail Making Test

WCST: Wisconsin Card Sorting Test

Y-BOCS: Yale-Brown Obsessive Compulsive Scale

# INTRODUCTION

Obsessive–compulsive disorder (OCD) is a relatively common and chronically disabling neuropsychiatric disorder with an average lifetime prevalence of 2%–3% (1). The OCD is characterized by intrusive thoughts which the patient attempts to neutralize by either mental or motoric compulsive acts (2). And the typical themes of these intrusive thoughts occurring in OCD relate to contamination fear, symmetry, aggression, sexuality and religious ideas (3). Cognitive behavioral models of OCD pathophysiology posit dysfunctional inhibitory control system during symptom provocation paradigms and neuropsychological tests requiring executive control of motor inhibition and cognitive flexibility (4).

Taking heterogeneous symptomatology of the OCD into consideration, more common disease characteristics encompassing diverse OC symptom dimensions are needed. Endophenotypes are measurable disease-associated traits which could occur even during the subclinical disease status and have a simpler relationship with underlying genes than clinical measurements, assisting in the identification of genes that result in vulnerability to the disorder (5). From this perspective, neurocognitive dysfunctions are considered among the most promising candidates for endophenotypes in many psychiatric disorders including OCD (6), in which executive dysfunctions in the domains of nonverbal memory (7), cognitive flexibility (8) and response inhibition are among the most frequently replicated findings. Executive dysfunctions may qualify as a suitable endophenotype candidate for OCD. Concordance rates in neuropsychological task performance suggest that decision-making and planning deficits aggregate in these families and therefore might be a heritable component of OCD (9).

The neuro-anatomical model of obsessive-compulsive disorder (OCD) proposes that the clinical symptoms and related cognitive deficits of OCD are caused by dysfunctional cortico–striato–thalamo-cortical (CSTC) loops. The CSTC loops are composed of several parallel running loops encompassing prefrontal areas of the orbitofrontal cortex (OFC), the anterior cingulate gyrus (ACC), the striatum including caudate nucleus and thalamus (10). Dysfunctional CSTC loops could result in primary executive dysfunction, which may affect secondary nonverbal memory function by way of impaired information processing strategies (11). And performances of the Rey–Osterrieth Complex Figure Test (RCFT) – a neuropsychological test assessing visuospatial constructional ability, visual memory, and executive function of visual organizational strategies (12) - and the Trail Making Test (TMT) – a neuropsychological test measuring attention, visuospatial working memory and executive function of the set-shifting (13) – are shown to be significantly impaired in OCD patients(13). But the exact neural correlates of these neuropsychological impairments have not been investigated in OCD yet.

As an integrative center of several cognitive-behavioral, emotional-autonomic and motor neural networks (14, 15), the anterior cingulate cortex (ACC), one of the prefrontal targets of the CBTC loops, has been suggested as a putative neural correlate of executive dysfunction in OCD. ACC is anatomically, metabolically and functionally altered in OCD. ACC is linked to difficulties with set-shifting (16), decision making and both action (17) and error-monitoring (18, 19) in OCD patients. Previous structural neuroimaging studies have revealed some structural alterations in ACC morphology - namely, reduced grey matter (GM) volume of the left ACC (20), left caudal ACC (21) and bilateral dorsal ACC (22) – in the OCD patients, when compared to the healthy control (HC) group. ACC-associated white matter (WM) alterations were also reported, as white

matter (WM) volume (23) and fractional anisotropy (24) of the left ACC were significantly reduced in the OCD group.

Not only these structural changes, in OCD patients, ventral frontal-striatal areas such as OFC, caudate head, insula and ACC are hyper-activated during symptom provocation (25) or during resting states (26). And in several task-based fMRI studies, dysfunctional ACC activation in OCD has been linked to the executive dysfunction of set-shifting(16), action monitoring (17), error-processing (18) and cognitive control of emotion (27). Also, in a recent clinical study using Tower of London test as a task, as task load increases, pediatric drug-naïve OCD patients showed more recruitment of ventro-lateral and medial prefrontal cortex and insula and ACC when compared to HC group, which were normalized after cognitive behavioral treatment (CBT) sessions (28). Using EEG source localization method, hyperactive error-related negativities (ERNs), which is related to the obsessive thoughts and compulsive acts of OCD patients and is thought to reflect the activities of a hyperactive cortico-striatal circuit during action monitoring, was detected in the rostral ACC of OCD patients with severe OC symptomatology (29). Likewise, previous <sup>1</sup>H-magnetic resonance spectroscopy (MRS) studies have reported reduced N-acetyl-aspartate (NAA) level and glutamate-glutamine level in the ACC of OCD patients (30, 31). All these anatomical and metabolic alterations and changes in anatomical connectivity between ACC and other brain area are reflected in brain studies using EEG and functional magnetic resonance imaging (fMRI). In an EEG study for severe OCD patients, error-related negativities (ERNs) were localized to the rostral ACC (29). (32, 33). But to overcome the limitations of previous task-based fMRI studies, in which the exact loci and extents of ACC activation became diverse depending upon the function explored and the task employed (34), another functional neuroimaging approach which could demonstrate the pattern of ACC



functioning in wider dimension of .cognitive function has been required.

As both cognitive functioning and OC symptomatology are manifested over the integrated activity of multiple brain regions, investigating the coupling between several brain regions could be more useful to understand complex brain functioning. These relationships between brain regions can be measured under the notion of 'functional connectivity', the spatiotemporal correlations between spatially distinct brain regions (35). And among the functional connectivities, the resting-state functional connectivity networks (rs-FCNs), spontaneous low-frequency fluctuations (0.01–0.08 Hz) of resting-state blood oxygenation level-dependent signal, have been widely used to characterize the intrinsic or spontaneous neuronal activity of the normal subjects as well as the psychiatric or neurologic patient groups (36). The rs-FCN approach could reveal the degree of functional-metabolic coherence between brain regions during task-free status (37), in which diverse rs-FCNs including the default mode network (DMN) – related to internally oriented cognition and deactivated when required to act goal-directed behavior (38), the dorsal attention network (39), the somatosensory network (40) and the paralimbic network(41) coexist along the whole brain area.

The ACC, composed of cyto-architectonically (42) and functionally (14) heterogeneous sub-regions, works as the integrative centers for several cognitive–behavioral, emotional–autonomic, and motor neural networks. ACC is structurally (42) and functionally (14) heterogeneous. The existence of motor, cognitive and affective subdivisions of ACC in healthy subjects has been revealed in previous task-based neuroimaging studies (14). And with the use of the rs-FCN approach, the presence of rostral ACC-based affective network and caudal ACC-based fronto-parietal attention networks and reciprocal relationship between them was already elucidated in healthy subjects (43).

In case of OCD, the functional heterogeneity of ACC, which has been demonstrated in the multiple task-based fMRI studies with contradictory findings (44), has not yet been investigated using the unique rs-FCNs methodology. For ACC-centered connectivity in OCD, one of our recent study found out weakened rs-FCN between the posterior cingulate cortex (PCC) and subcallosal ACC (Brodmann area (BA) 24) in DMN (45), the clinical importance of which not elucidated yet. In addition, not only the weakened rs-FCN strength between dorsal ACC (BA 24) and right OFC in adult OCD patients (46) but also the decreased rs-FCN strength between rostral ACC and caudate head in child OCD patients (mean age 11.0 years old) (47) showed correlations with higher Yale-Brown Obsessive Compulsive Scale (Y-BOCS) scores. But rs-FCN approaches for OCD with regard to the cytoarchitectonic/functional sub-regions of ACC as well as their cognitive implications are rare.

In this study, we used rs-FCN approach using MRI to investigate changes in topological organization in intrinsic or spontaneous brain activities in top-down control networks in patients with OCD. <sup>w</sup>e separately examined the FCs of five selected seed regions of interest (ROIs) systematically placed throughout the ACC. And we also made all subjects to conduct five neuropsychological tests measuring diverse dimensions of the executive function. Finally, we analyzed the correlations between the derangements of ACC-seeded rs-FCN subdivision versus neuropsychological test performances. In this study, we hypothesized that the ACC seed-based rs-FCN would show disrupted strength in OCD. This altered strength of functional connectivity could be a reflection of inefficient information processing in OCD, which might correlate with the performance level of neuropsychological tests for executive function measurement.

# MATERIALS AND METHODS

## 1. Subjects

All subjects included in this study had participated in our previous functional magnetic resonance imaging (fMRI) study using AFNI software (<http://afni.nimh.nih.gov/afni>) (45), from which one OCD patient was not involved in this study because of technical restraints in fMRI image preprocessing using Statistical Parametric Mapping 8 (SPM8) software (<http://www.fil.ion.ucl.ac.uk/spm>). Twenty-one patients (15 men and six women) who fulfilled the Diagnostic and Statistical Manual of Mental Disorders, 4<sup>th</sup> edition (DSM-IV) criteria for OCD, as diagnosed using the Structured Clinical Interview for DSM-IV (SCID-IV), were recruited from the OCD clinic at Seoul National University Hospital. Fifteen patients were drug-naïve, and the remaining six had been drug-free for at least 4 weeks at inclusion. Twenty-two gender- and age-matched healthy controls were also recruited. The SCID Non-patient Version (SCID-NP) was used to assess psychiatric disorders in controls. Exclusion criteria included a lifetime history of major psychiatric disorders, significant head injury, seizure disorder, or mental retardation. The current study was approved by the Institutional Review Board at Seoul National University Hospital, and written informed consent was obtained from all subjects after the procedures had been fully explained.

## 2. Clinical and Neuropsychological Assessments

Obsessive-compulsive symptom severity was measured with the 10-item clinician-rated Yale–Brown Obsessive Compulsive Scale (Y-BOCS) (48). The Beck Depression Inventory (BDI) (49) and the Beck Anxiety Inventory (BAI) (50) were also administered to measure the severity of depression

and anxiety, respectively. The Korean version of the Wechsler Adult Intelligence Scale (K-WAIS) (51) was administered to measure the IQ.

To measure the degrees of the executive dysfunction which might be the manifestation of dysfunctional CBTC loop in the OCD (11), we used 5 neuropsychological tests (52, 53) in this study: the Wisconsin Card Sorting Test (WCST), the Controlled Oral Word Association Test (COWAT), the Stroop Color-Word Test (SCWT), Part A and B of the TMT and the RCFT.

The WCST measures the executive function of abstract reasoning, concept formation and attentional set shifting in clinical populations including the OCD patients (54). In this study, we used the conventional WCST form (55) for test administration and scoring. In the Heaton's version, subjects are required to find the correct classification principle for color, form or number cards by way of trial, error and examiner feedback. Once the subject chooses the correct rule, they must maintain this sorting principle (or set) across changing stimulus conditions. Then the classification principle changes after ten consecutive correct responses without warning to the subjects, and the sorting process continues until all cards are sorted or a maximum of six correct sorting criteria have been reached. During these process of card sorting, perseverative responses of sorting the cards consecutively in the same way or repeating the previous principle (56), is selected in this study as a proper measure of so called 'frontal executive functioning' (57).

The COWAT performance reveals the degree of verbal fluency, which requires properly organized verbal retrieval, ongoing monitoring of the verbal search set in working memory and inhibition of inappropriate responses (58). We used 'Category Fluency' score as a representative score of cognitive flexibility in this study.

The SCWT color-word page provides conflicting visual stimuli to measure

the executive function of selective attention and response inhibition, which is well reflected in reaction time (59).

The TMT (60) from the Halstead–Reitan Neuropsychological Test Battery is a measure of attention and executive function. Drawing on worksheets, individuals must first connect consecutively numbered circles (Part A) and then connect consecutively numbered and lettered circles by alternating the two sequences (Part B). The TMT Part A reflects attention capacity, concentration and psychomotor speed of the subjects, and the TMT part B requires subjects of intact executive function named set-shifting (13).

The RCFT, originally designed by Rey (61) and scored by Osterrieth (62), is one of the neuropsychological tests which has been widely used to evaluate visuospatial constructional ability and visual (non-verbal) memory functions in the clinical population (63). During the procedure of RCFT performance, subjects were instructed to copy the RCFT (64) figure and then draw what they remembered immediately. And for the comprehensive scoring of the RCFT performances in terms of visuo-constructional ability as well as the executive functioning, we used the Boston Qualitative Scoring System (BQSS) (65), which provided us with the 17 kinds of qualitative scores and 6 summary scores. In this study, we have chosen the Organization Summary Score, which is the arithmetic sum of the copy condition Fragmentation score (measure of information integration) and Planning score (reflecting the overall planning ability based on the order, placement and overall integrity of the depicted elements), as a representative of the executive function measurement (63).

### **3. fMRI Image Acquisition**

Functional images were acquired using a 1.5 T MAGNETOM Avanto

scanner (Siemens, Erlangen, Germany). For the resting state with no task performances, functional scans for the whole brain were acquired in 25 contiguous axial slices approximately parallel to the anterior-posterior commissure plane with interleaved multi-slice echo-planar imaging according to the following parameters: TR = 2.34 s, TE = 52 ms, field of view = 22 cm, flip angle = 90°, voxel size = 3.44 × 3.44 × 5 mm, slice thickness = 5 mm, no inter-slice gap.

The resting state fMRI scanning procedure consisted of 120 volumes during 4.68 min. During these resting state experiments, subjects were lying in the dark with their eyes closed, with instructions to relax as much as possible and think about nothing in particular. Meanwhile, the spontaneous brain activity level of each subject was measured throughout the experimental period. Additionally, T1-weighted high-resolution structural images, obtained using a magnetization-prepared rapid acquisition gradient echo sequence, were acquired in 176 contiguous axial slices to co-register and normalize the echo-planar images to the Montreal Neurologic Institute (MNI) template. Imaging parameters for these structural images were as follows: TR = 1.16 s, TE = 4.76 ms, field of view = 23 cm, flip angle = 15°, voxel size = 0.45 × 0.45 × 0.90 mm, slice thickness = 0.9 mm, no inter-slice gap.

## **4. fMRI Preprocessing**

Functional data were preprocessed and analyzed using SPM8 (<http://www.fil.ion.ucl.ac.uk/spm>).. The first four volumes of the functional images were removed to eliminate the non-equilibrium effects of magnetization. Preprocessing steps included slice-timing correction for interleaved acquisition, head motion correction, spatial normalization into

standard stereotactic MNI spaces with re-sampling to 3-mm cubic voxels, and spatial smoothing using a Gaussian kernel of 6-mm full width at half maximum.

The following additional preprocessing steps were performed for the resting state functional connectivity analysis (66). The resting-state fMRI Data Analysis Toolkit (REST version 1.5, <http://resting-fmri.sourceforge.net>) was then used to remove the linear trend of time courses and for temporal band-pass filtering ( $0.01\text{Hz} < f < 0.08\text{Hz}$ ), as a major research interest with regard to the resting-state fMRI is in slow-changing temporal activation. Before the correlation analysis, signals from the regions of no interest and the first temporal derivatives (head motion parameters, global, white matter, and cerebrospinal fluid signals) were regressed out to reduce spurious correlations derived from cardiac and respiratory fluctuations (67, 68).

## **5. Functional Connectivity: Seed Generation**

We examined the FCs of five selected seed regions of interest (ROIs) out of 16 seed ROIs systematically placed throughout the ACC in two arrays designated superior (S) and inferior (I), following the methods outlined in other study (43). These five ACC seeds selected in the present study refer to the samples of five domains of the self-regulatory functions associated with the ACC; namely, motor control, attention/cognitive control, conflict monitoring, mentalizing, and emotional regulation (69). Seed S1 (MNI coordinates:  $x = 5, y = -10, z = 47$ ), located in the paracentral lobule and corresponding to BA 31, is in the caudal ACC. The dorsal ACC (dACC), the location of Seed S3 ( $x = 5, y = 14, z = 42$ ) is in the BA 32. Seed S5 ( $x = 5, y = 34, z = 28$ ) is located in the rostral section of the supragenual ACC. Seed S7 ( $x = 5, y = 47, z = 11$ ), which resides in the medial frontal gyrus of BA 32,

corresponds to the perigenual ACC. Finally, Seed I9 ( $x = 5$ ,  $y = 25$ ,  $z = -10$ ) is located in the subgenual ACC, corresponding to BA 32.

## **6. Functional Connectivity Map Generation**

Using the “functional connectivity” function of REST, we examined rs-FC maps of the five predefined ACC seed ROIs for each subject. Each spherical seed had a radius of 3 mm in a 3 x 3 x 3 mm voxel-sized space. For each participant, we calculated the mean time series of each seed ROI by averaging across all voxels within the seed. A correlation map for each seed was produced by computing the correlation coefficients voxel-wise. Finally, the correlation coefficient maps were converted into z maps by Fisher’s z transform to improve normality.

## **7. Statistical Analysis**

The demographic and clinical data for the OCD and control groups were compared with independent *t*-tests for continuous variables and chi-square tests for categorical variables using PASW 18.0 (SPSS Inc., Chicago, IL, USA).

To reveal within-group functional connectivity patterns, we entered each single-subject intrinsic network component into a voxel-wise one-sample *t*-test in SPM8 to create statistical z maps of the intrinsic networks based on each of the five ACC seeds (uncorrected  $p < 0.001$ ,  $k > 10$ ). After obtaining the regions constituting the intrinsic organization for each group, a binary mask was created using the ImCalc toolbox in SPM8, separately for both groups. Then, a combined mask was generated by combining these two masks.



Differences in the rs-FCNs between the OCD group and the control group were analyzed using two-sample *t*-tests, with a previously described masking procedure, to avoid detection of clusters that did not appear in the ACC network map. The resulting statistical maps are presented according to the threshold of uncorrected  $p < 0.001$  and a cluster threshold of  $k > 10$  for multiple comparisons, which corresponds to the corrected threshold of  $p < 0.05$ , as determined by AlphaSim in AFNI (<http://afni.nih.gov/afni/docpdf/AlphaSim.pdf>).

Correlation coefficients (Z scores) of specific clusters with significantly different rs-FC strengths between the two groups were extracted using the MarsBar toolbox (<http://marsbar.sourceforge.net>) to test the relationship between rs-FC strength versus neuropsychological test performance and Y-BOCS scores (total score, obsession subscale and compulsion subscale) in the OCD group. Pearson's simple correlation coefficient was used to investigate the associations between these five ACC seed-related rs-FC strengths versus several neuropsychological test sub-scores measuring executive functioning and Y-BOCS scores. To fully explore the possibility of the ACC-based rs-FCNs as a biomarker reflecting executive dysfunction and obsessive-compulsive symptomatology in OCD patients, we admitted a more exploratory threshold of  $p < 0.05$  (uncorrected, two-tailed) rather than the strict threshold drawn by the Bonferroni–Holms procedure. And post-hoc analyses to examine the impact of some clinical variables of BDI and BAI scores rs-FCN alteration were conducted, as these two variables showed significant differences between the OCD group and healthy controls.

# RESULTS

## 1. Demographic and Clinical Characteristics

The demographic and clinical characteristics of the subjects are presented in Table 1 and 2. We found no significant differences between the groups regarding age, gender, years of education and IQ. In contrast, significant differences were observed between the OCD group and the control group for the BAI and BDI scores ( $p < 0.05$ , two-tailed)

In this study, no significant differences were observed in RCFT organization score and COWAT category fluency score. In contrast, the OCD group and control group showed significant differences in the the WCST perseverative response score, the SCWT reaction time, and reaction time for TMT part A/B ( $p < 0.05$ , two-tailed).

**Table 1.** Demographic and clinical characteristics of OCD group and healthy control

	OCD (n=21)	Healthy control (n=22)	t-score	p-value
<b><i>Demographic characteristics</i></b>				
Age(year)	25.10±7.13	24.36±4.02	0.412	0.683
Gender(M/F)	15/6	16/6	-	-
Education(year)	14.38±3.81	14.41±1.62	-0.031	0.975
Estimated IQ	108.57±13.12	112.50±13.05	-0.961	0.343
Illness duration (year)	7.39±5.45	-	-	-
<b><i>Clinical rating scales</i></b>				
Y-BOCS score				
Obsession	11.48±4.52	-	-	-
Compulsion	10.33±4.59	-	-	-
Total score	21.81±6.96	-	-	-
BAI score	18.57±15.02	4.05±5.20	4.177	<0.001*
BDI score	16.52±11.01	3.65±4.93	4.870	<0.001*

*Abbreviations:* OCD: obsessive-compulsive disorder; Y-BOCS: Yale-Brown Obsessive Compulsive Scale; BAI: Beck Anxiety Inventory; BDI: Beck Depression Inventory.

Data are given as mean ± standard deviation.

\*  $p < 0.05$

**Table 2.** Neuropsychological test scores of the subjects


	OCD (n=21)	Healthy control (n=22)	t-score	p-value
WCST Perseverative response	9.05±4.02	6.45±3.07	2.386	0.022*
COWAT category fluency	38.81±8.94	41.41±9.26	-0.936	0.355
SCWT RT	109.33±21.71	89.23±26.52	2.713	0.010*
TMT-B RT	75.10±31.94	56.05±17.17	2.452	0.019*
TMT-A RT	33.10±12.76	24.14±7.19	2.77	0.010*
RCFT organization score	6.62±1.24	6.86±1.17	-0.665	0.510

Data are given as mean ± standard deviation. \* $p < 0.05$


*Abbreviations:* OCD: Obsessive-Compulsive Disorder; WCST: Wisconsin Card Sorting Test; COWAT: Controlled Oral Word Association Test; SCWT: Stroop Color Word Test; TMT: Trail Making Test; RCFT: Rey-Osterrieth Complex Figure Test

## 2. Resting State Functional Connectivity


Overall, both the OCD and control groups exhibited a distinct segregation of ACC in rs-FCNs along the rostro-caudal axis, which matched with previous studies on functional differentiation within the ACC (43, 69). The specific intra-group and inter-group findings for each of the ACC seed regions are described below. And rs-FCN maps of the OCD group for each five of ROI seeds are presented as separated figures (Figure 1 for S1 seed; Figure 2 for S3 seed; Figure 3 for S5 seed; Figure 4 for S7 seed; and Figure 5 for I9 seed).




**Figure 1.** Positive (depicted with red) and negative (depicted with blue) resting state functional connectivity networks between the S1 ACC seed region and other brain area in Obsessive-Compulsive Disorder group. Results are displayed at  $P < .001$ , uncorrected.



**Figure 2.** Positive (depicted with red) and negative (depicted with blue) resting state functional connectivity networks between the S3 ACC seed region and other brain area in Obsessive-Compulsive Disorder group. Results are displayed at  $P < .001$ , uncorrected.




**Figure 3.** Positive (depicted with red) and negative (depicted with blue) resting state functional connectivity networks between the S5 ACC seed region and other brain area in Obsessive-Compulsive Disorder group. Results are displayed at  $P < .001$ , uncorrected.



**Figure 4.** Positive (depicted with red) and negative (depicted with blue) resting state functional connectivity networks between the S7 ACC seed region and other brain area in Obsessive-Compulsive Disorder group. Results are displayed at  $P < .001$ , uncorrected.





**Figure 5.** Positive (depicted with red) and negative (depicted with blue) resting state functional connectivity networks between the I9 ACC seed region and other brain area in Obsessive-Compulsive Disorder group. Results are displayed at  $P < .001$ , uncorrected.

## **2.1. Five ACC Seed-Related rs-FCNs: Overlapping Findings of positive correlation**

Caudal Seed S1 in the patients with OCD and healthy controls correlated with a network of premotor and supplementary motor areas (BA 6), the primary somatosensory cortex (BA 3), the dorsal ACC (BA 24/32), the primary motor cortex (BA 4), the pars opercularis (BA 44), the insula (BA 13), the posterior cingulate cortex (PCC) (BA 31), the claustrum, and the paracentral lobule (BA 5) (Figure 1 and 6).


Seed S3, which resides in the dorsal ACC, demonstrated significant rs-FC in the caudal ACC (BA 24/32), PCC (BA 31), premotor and supplementary motor area, precuneus (BA 7), postcentral gyrus (BA 7), primary motor cortex, primary somatosensory cortex (BA 1/2/3), orbitofrontal cortex (OFC) (BA 10/47), superior temporal gyrus (BA 22), claustrum, supramarginal gyrus of the inferior parietal lobule (IPL) (BA 40), Broca's area (BA 44/45), insula, primary auditory cortex (BA 42), putamen, and dorsolateral prefrontal cortex (DLPFC) (BA 9) in both groups (Figure 2 and 6). These results are in accordance with the previous functional connectivity findings (70).

The rostral ACC seed region S5 in both groups showed significant rs-FC with the OFC (BA 10/11/ 47), DLPFC (BA 8/9/46), premotor area, Broca's area, caudal/dorsal/perigenual ACC (BA 32), PCC (BA 23/31), caudate body, putamen, claustrum, insula, precuneus (BA 7), supramarginal gyrus of the IPL, superior occipital gyrus (BA 19), and tonsil/tuber of the cerebellum (Figure 3 and 6).

Perigenual ACC Seed S7 in both groups revealed rs-FC with the OFC (BA 10/47), DLPFC (BA 8/9), pregenual (BA 33)/ventral (BA 24)/perigenial (BA 32) ACC, premotor cortex, subcallosal gyrus (BA 34), precuneus (BA

7/19/31), PCC (BA 23/31), superior parietal lobule (BA 7), angular (BA 39) and supramarginal gyrus (BA 40) of the IPL, Wernicke's area (BA 22), temporal pole (BA 38), fusiform gyrus (BA 20), insula, middle (BA 21) and inferior (BA 20/21) temporal cortices, head/body of the caudate, thalamus, and culmen/pyramis/inferior semi-lunar lobule of the cerebellum (Figure 4 and 6).


Seed I9, located in the subgenual ACC, was associated with an extensive pattern of correlated activity in the OFC (BA 10/11), pars opercularis, perigenual (BA 32) /subgenual (BA 25) ACC, subcallosal gyrus, middle (BA 19) and inferior (BA 20) temporal cortices, uncus (BA 34/38), caudate head, tonsil and pyramis of the cerebellum in both groups (Figure 5 and 6).



**Figure 6.** Positive resting state functional connectivity networks of the 5 ACC seed ROIs in OCD and healthy control. Significant within-group functional connectivity maps of the 5 ACC seeds are placed on 3-dimensional anatomical images. Results are displayed at  $P < .001$ , uncorrected.

## **2.2. Five ACC Seed-Related rs-FCNs: Between-Group Analyses of positive correlation**

Between-group analyses revealed that the rs-FC between the ACC S1 seed and the left claustrum, between the ACC S3 seed and the left pulvinar thalamus, between the ACC S5 seed and the right claustrum, and between the ACC S7 seed and the right IPL were significantly weaker in the OCD group. In contrast, the rs-FC between the ACC I9 seed and the right OFC was significantly stronger in the OCD group ( $p < 0.001$ ,  $k > 10$ ) (Figure 7, Table 3). When additional multiple regression analyses (forced entry method) which put forward BDI, BAI scores as well as the 'diagnosis' as predictor variables were performed, no predictor variables other than the 'diagnosis' survived the statistical significance threshold of  $p < .05$  (Table 4).



**Figure 7.** Significant between-group (main effect) differences in positive rs-FCNs of the 5 ACC seed ROIs. The results are placed on (A) 3-dimensional, (B) 2-dimensional and (C) anatomical brain images. Colored overlays correspond regions demonstrating weaker functional connectivity in OCD patients, with respect to red (S1), green (S3), blue (S5) and pink (S7); yellow, regions depicting stronger functional connectivity with the respective ACC I9 seed region in patients vs controls. Results are displayed at  $P < .001$ , uncorrected.

**Table 3.** Regional difference in resting state functional connectivity network between the patients with OCD and healthy controls

brain region	laterality	BA	Peak coordinates (MNI)			T score	Z score	p value	cluster size
			X	y	z				
<b><i>S1_R: control&gt;OCD</i></b>									
Claustrium	L	-	-39	-24	0	4.24	3.84	<0.001	29
<b><i>S3_R: control&gt;OCD</i></b>									
thalamus, pulvinar	L	-	-15	-24	15	4.87	4.3	<0.001	21
<b><i>S5_R: control&gt;OCD</i></b>									
Claustrium	R	-	27	21	3	4.34	3.92	<0.001	12
<b><i>S7_R: control&gt;OCD</i></b>									
Supramarginal gyrus	R	40	60	-48	48	4.49	4.02	<0.001	11
<b><i>I9_R: OCD&gt;control</i></b>									
Frontopolar cortex	R	10	18	66	9	3.81	3.5	<0.001	11

*Abbreviations:* BA: Brodmann area; MNI: Montreal Neurological Institute; OCD: Obsessive-compulsive disorder; L: left; R: right.

The thresholds of statistical significance in between-group analyses were set to  $p < 0.001$  (uncorrected) and  $k$  (voxel threshold)  $> 10$ .

**Table 4.** Significance of diagnosis in predicting strength of positive resting state functional connectivity\*

	$R^2$ (p value)	Predictor variables	B	SE B	t score	p value
<b>S1 vs. Lt. claustrum</b>	0.438	Diagnosis	-0.168	0.054	-3.077	0.004†
	(p<.001†)	BDI	-0.002	0.003	-0.456	0.651
		BAI	-0.002	0.003	-0.930	0.358
<b>S3 vs. Lt. pulvinar</b>	0.387	Diagnosis	-0.218	0.068	-3.224	0.003†
	(p<.001†)	BDI	-0.007	0.004	-1.628	0.112
		BAI	0.004	0.003	1.313	0.197
<b>S5 vs. Rt. Claustrum</b>	0.411	Diagnosis	-0.253	0.054	-4.701	<0.001†
	(p<.001†)	BDI	0.005	0.003	1.541	0.132
		BAI	-0.001	0.003	-0.519	0.607
<b>S7 vs. Rt. supramarginal gyrus</b>	0.355	Diagnosis	-0.242	0.083	-2.900	0.006†
	(p=.001†)	BDI	0.003	0.005	0.660	0.513
		BAI	-0.006	0.004	-1.482	0.146
<b>I9 vs. Rt. frontopolar cortex</b>	0.274	Diagnosis	0.329	0.095	3.446	0.001†
	(p=.005†)	BDI	-0.006	0.006	-1.084	0.285
		BAI	0.002	0.004	0.357	0.723

\*Results were obtained from multiple regression analyses also including BDI/BAI score as predictors. †p< .05.

*Abbreviations:* BAI= Beck Anxiety Inventory; BDI= Beck Depression Inventory; Lt.= left; Rt.= right;  $R^2$ = coefficient of determination; SE= standard error



### **2.3. Five ACC Seed-Related rs-FCNs: negative correlation**

There were also marked distinctions in negative relationships (“anticorrelations”) between ACC and other brain regions, with systematic variations along the axis of rostral to caudal portions of ACC.

The subgenual seed I9, and superior rostral seed S7 negatively predicted activity within a number of posterior brain regions linked to sensorimotor and attentional domains, including superior, inferior and posterior parietal regions, temporal cortices, portions of extrastriate and striate visual cortices, and superior cerebellar regions for both groups of OCD and HC, with no significant differences in the rs-FC anti-correlation strength between these two groups (Table 5).

Centrally located superior seed s5 negatively predicted activity within portions of the occipito-temporal lobes and superior, posterior and parietal lobes in the HC group. In contrast, these patterns of anti-correlations regarding S5 seed were less distinctive in the OCD group, especially for the brain regions of right fusiform gyrus (BA 18), right associative visual cortex (BA 19) and left primary somatosensory cortex (BA 2) (uncorrected  $p < .001$ ,  $k > 10$ ) (Table 5).

In case of caudally located ACC seeds of s1 and s3, negatively predicted activity was extended to the whole areas of frontal lobe - many of the same regions positively predicted by extreme rostral seeds – as well as in the posterior cerebral cortices. However, inter-group differences in the strength of anti-correlation were located in the temporo-occipital and cerebellar cortices, as OCD group showed more attenuated rs-FC anti-correlation between s3 seed versus right parahippocampal gyrus (BA 19), right lingual gyrus (BA 18), left associative visual cortex (BA 19) and cerebellum (right culmen). For the extremely caudal-located seed s1, significantly attenuated

rs-FC anti-correlation in the OCD group was not found (uncorrected  $p < .001$ ,  $k > 10$ ) (Table 5).

When additional multiple regression analyses (forced entry method) which put forward BDI, BAI scores as well as the 'diagnosis' as predictor variables were performed, no predictor variables other than the 'diagnosis' survived the statistical significance threshold of  $p < .05$  (Table 6).

**Table 5.** Regional difference in negative correlations of resting state functional connectivity network between the patients with obsessive-compulsive disorder and healthy controls

brain region	laterality	BA	Peak coordinates (MNI)			T score	Z score	p value	cluster size
			x	y	z				
<b>S1_R: NS</b>									
<b>S3_R: control&gt;OCD</b>									
Culmen of cerebellum	R	-	18	-42	-9	4.39	3.95	<0.001	36
Parahippocampal gyrus	R	19	21	-48	0	3.85	3.54	<0.001	
Associative visual cortex	L	19	-27	-87	9	4	3.66	<0.001	15
Lingual gyrus	R	18	24	-75	-6	3.61	3.34	<0.001	11
<b>S5_R: control&gt;OCD</b>									
Fusiform gyrus	R	18	24	-87	-15	4.44	3.99	<0.001	11
Associative visual cortex	R	19	39	-78	24	4.39	3.95	<0.001	14
Primary somatosensory cortex	L	2	-45	-33	63	4.37	3.93	<0.001	28
<b>S7_R &amp; I9_R: NS</b>									

*Abbreviations:* BA: Brodmann area; MNI: Montreal Neurological Institute; OCD: Obsessive-compulsive disorder; L: left; R: right.

The thresholds of statistical significance in between-group analyses were set to  $p < 0.001$  (uncorrected) and  $k$  (voxel threshold)  $> 10$ .

**Table 6.** Significance of diagnosis in predicting strength of negative resting state functional connectivity\*

	$R^2$ (p value)	Predictor variables	B	SE B	t score	p value
<b>S3 vs. Rt. Parahippocampal gyrus</b>	0.356	Diagnosis	0.325	0.075	4.317	<0.001 <sup>†</sup>
	(p=.001 <sup>†</sup> )	BDI	-0.004	0.005	-0.883	0.382
		BAI	-0.001	0.004	-0.286	0.776
<b>S3 vs. Lt. associative visual cortex</b>	0.276	Diagnosis	0.278	0.082	3.385	0.002 <sup>†</sup>
	(p=.005 <sup>†</sup> )	BDI	-0.004	0.005	-0.726	0.472
		BAI	0.001	0.004	0.161	0.873
<b>S3 vs. Rt. Lingual gyrus</b>	0.290	Diagnosis	0.282	0.078	3.617	<0.001 <sup>†</sup>
	(p=.004 <sup>†</sup> )	BDI	-0.006	0.005	-1.187	0.243
		BAI	0.001	0.004	0.357	0.723
<b>S5 vs. Rt. Fusiform gyrus</b>	0.388	Diagnosis	0.302	0.065	4.640	<0.001 <sup>†</sup>
	(p<.001 <sup>†</sup> )	BDI	-0.007	0.004	-1.805	0.079
		BAI	0.001	0.003	0.452	0.654
<b>S5 vs. Rt. Visual associative cortex</b>	0.357	Diagnosis	0.245	0.075	3.279	0.002 <sup>†</sup>
	(p=.001 <sup>†</sup> )	BDI	-0.006	0.005	-1.210	0.233
		BAI	0.005	0.003	1.571	0.124
<b>S5 vs. Lt. primary somatosensory cortex</b>	0.383	Diagnosis	0.179	0.064	2.789	0.008 <sup>†</sup>
	(p<.001 <sup>†</sup> )	BDI	-0.001	0.004	-0.297	0.768
		BAI	0.004	0.003	1.497	0.142

Results were obtained from multiple regression analyses also including BDI/BAI score as predictors. <sup>†</sup>p< .05

*Abbreviations:* BAI= Beck Anxiety Inventory; BDI= Beck Depression Inventory; Lt.= left; Rt.= right;  $R^2$ = coefficient of determination; SE= standard error


### **2.3. Correlations between altered rs-FCNs versus Y-BOCS scores**

For the regions showing significant differences in functional connectivity strength between OCD patients and controls, increased positive rs-FC strength between ACC Seed S5 and the right claustrum correlated with higher Y-BOCS total score ( $r=.548$ ,  $p=.010$ ) and Y-BOCS Compulsion subscale ( $r = 0.497$ ,  $p = 0.022$ ) in the OCD group. Regarding the Y-BOCS Obsession subscale, stronger positive rs-FC between the ACC S1 seed versus left claustrum could predict lower Y-BOCS Obsession subscale in OCD patients ( $r=-.487$ ,  $p=.025$ ) (Figure 8, Table 7). In terms of the negative rs-FCNs, stronger negative rs-FC between the ACC S3 seed versus right parahippocampal gyrus showed correlations with lesser compulsive symptomatology measured using the Y-BOCS Compulsion subscale ( $r=.481$ ,  $p=.027$ ) (Figure 9, Table 7). No other statistically significant correlations between ACC seed-based regional rs-FC strength versus Y-BOCS scores were found ( $p < 0.02$ , two-tailed).


**Table 7.** Correlation between altered regional resting state functional connectivity strength versus Y-BOCS scores in Obsessive-Compulsive disorder subjects

rs-FCN		Obsession	Compulsion	Total
<b>Positive correlation</b>				
<b>S1 vs. Lt. claustrum</b>	Correlation	-0.487	0.123	-0.236
	Significance	0.025*	0.597	0.303
<b>S3 vs. Lt. pulvinar</b>	Correlation	0.186	-0.133	0.033
	Significance	0.420	0.565	0.887
<b>S5 vs. Rt. Claustrum</b>	Correlation	0.340	0.497	0.548
	Significance	0.132	0.022*	0.010*
<b>S7 vs. Rt. Supramarginal gyrus</b>	Correlation	-0.205	-0.066	-0.177
	Significance	0.373	0.776	0.444
<b>I9 vs. Rt. frontopolar cortex</b>	Correlation	-0.171	-0.065	-0.154
	Significance	0.459	0.780	0.506
<b>Negative connectivity</b>				
<b>S3 vs. Lt. associative visual cortex</b>	Correlation	-0.049	0.018	-0.019
	Significance	0.834	0.937	0.933
<b>S3 vs. Rt. Lingual gyrus</b>	Correlation	-0.046	0.171	0.083
	Significance	0.845	0.459	0.721
<b>S3 vs. Rt. parahippocampal gyrus</b>	Correlation	-0.291	0.481	0.128
	Significance	0.200	0.027*	0.581
<b>S5 vs. Lt. primary somatosensory cortex</b>	Correlation	0.294	-0.05	0.158
	Significance	0.196	0.831	0.493
<b>S5 vs. Rt. Fusiform gyrus</b>	Correlation	-0.369	0.044	-0.21
	Significance	0.100	0.849	0.360
<b>S5 vs. Rt. Associative visual cortex</b>	Correlation	0.137	0.107	0.159
	Significance	0.554	0.644	0.490

\*p<.05. *Abbreviations:* Lt.= left; rs-FCN= resting state functional connectivity network; Rt.= right; Total= Y-BOCS total score



**Figure 8.** Correlation between positive regional functional connectivity strength and Y-BOCS scores in obsessive-compulsive disorder.



**Figure 9.** Correlation between negative regional functional connectivity strength and Y-BOCS scores in obsessive-compulsive disorder



## 2.4. Connectivity Correlations with Neuropsychological Measures

For the regions showing significant differences in functional connectivity strength between OCD patients and controls, increased rs-FC strength between ACC Seed S5 and the right claustrum correlated with higher RCFT organization scores in the OCD group ( $r = 0.541$ ,  $p = 0.011$ ) (Figure 10-(a), Table 8). Prolonged TMT-A reaction time was correlated with weaker rs-FC strength between ACC Seed S7 and the right supramarginal gyrus in the OCD group ( $r = -0.520$ ,  $p = 0.019$ ) (Figure 10-(b), Table 8). In addition, stronger positive rs-FC between the ACC I9 seed versus right frontopolar cortex also indicated shorter reaction time in TMT-A performance, with lower statistical significance ( $r = -.458$ ,  $p = .042$ ) than that of the ACC S7 seed. No other statistically significant correlations between ACC seed-based regional rs-FC strength versus neuropsychological test sub-scores were found ( $p < 0.05$ , two-tailed) (Table 8). In the healthy control group, lowered positive rs-FC between the ACC S7 seed versus right supramarginal gyrus indicated lesser WCST perseverative response ( $r = -.51$ ,  $p = .015$ ), better COWAT Category fluency performance ( $r = .477$ ,  $p = .025$ ) and the shorter TMT-B reaction time ( $r = -.576$ ,  $p = .005$ ). In addition, stronger positive rs-FC between the ACC S1 seed versus left claustrum indicated more perseverative response during WCST performance ( $r = .497$ ,  $p = .019$ ) (Table 9).

In the OCD group, negative rs-FCN alteration did not demonstrate statistically significant correlations with neuropsychological test performances (Table 10). In contrast, as negative rs-FC between the ACC S3 seed versus right lingual gyrus was weakened, the Category fluency score of COWAT became elevated in the healthy control group ( $r = .578$ ,  $p = .005$ ). ( $p < .05$ , two-tailed) (Table 11).

**Table 8.** Bivariate correlations between regional positive functional connectivity strength and neuropsychological test performance in Obsessive-Compulsive disorder patients (n=21)

Resting state functional connectivity network		WCST	COWAT	SCWT	TMT-B	TMT-A	RCFT
<b>S1 vs. Lt. claustrum</b>	Correlation	0.103	0.173	-0.166	0.093	-0.255	-0.136
	Significance	0.658	0.453	0.471	0.689	0.278	0.556
<b>S3 vs. Lt. pulvinar</b>	Correlation	-0.273	0.121	0.078	0.257	-0.151	0.089
	Significance	0.231	0.603	0.737	0.261	0.526	0.703
<b>S5 vs. Rt. Claustrum</b>	Correlation	0.094	-0.099	0.006	0.274	0.397	0.541
	Significance	0.684	0.670	0.981	0.230	0.083	0.011*
<b>S7 vs. Rt. supramarginal gyrus</b>	Correlation	-0.244	0.295	-0.238	-0.310	-0.520	0.086
	Significance	0.286	0.194	0.230	0.171	0.019*	0.710
<b>I9 vs. Rt. frontopolar cortex</b>	Correlation	0.187	0.332	0.008	-0.310	-0.458	0.005
	Significance	0.417	0.142	0.974	0.171	0.042*	0.983

*Abbreviation:* Correlation= Pearson's correlation coefficient; COWAT= Category fluency score of the Controlled oral word association test; Lt.= left; RCFT= Organization score of the Rey-Osterrieth complex figure test; Rt.= right; TMT-A= Reaction time for Trail-making test A; TMT-B= Reaction time for the Trail-making test B; SCWT= Reaction time for the color-word page of Stroop color word test; WCST= Perseverative response score of Wisconsin card sorting test. \*p<.05 (two-tailed)

**Table 9.** Bivariate correlations between regional positive functional connectivity strength and neuropsychological test performance in healthy controls (n=22)

Resting state functional connectivity network		WCST	COWAT	SCWT	TMT-B	TMT-A	RCFT
<b>S1 vs Lt. claustrum</b>	Correlation	0.497	-0.022	0.066	0.321	0.406	-0.016
	p value	0.019*	0.921	0.771	0.146	0.061	0.945
<b>S3 vs Lt. pulvinar</b>	Correlation	0.004	0.075	0.135	0.039	0.330	0.257
	p value	0.986	0.739	0.549	0.865	0.134	0.248
<b>S5 vs Rt. Claustrum</b>	Correlation	-0.173	0.026	-0.076	-0.214	-0.340	-0.119
	p value	0.441	0.910	0.737	0.339	0.122	0.599
<b>S7 vs Rt. supramarginal gyrus</b>	Correlation	-0.510	0.477	-0.084	-0.576	-0.407	0.116
	p value	0.015*	0.025*	0.710	0.005*	0.060	0.608
<b>I9 vs. Rt. frontopolar cortex</b>	Correlation	0.068	-0.225	0.121	0.223	0.402	0.372
	p value	0.765	0.315	0.591	0.319	0.063	0.089

*Abbreviation:* Correlation= Pearson's correlation coefficient; COWAT= Category fluency score of the Controlled oral word association test; Lt.= left; RCFT= Organization score of the Rey-Osterrieth complex figure test; Rt.= right; TMT-A= Reaction time for Trail-making test A; TMT-B= Reaction time for the Trail-making test B; SCWT= Reaction time for the color-word page of Stroop color word test; WCST= Perseverative response score of Wisconsin card sorting test. \*p<.05 (two-tailed)

**Table 10.** Bivariate correlations between regional negative functional connectivity strength and neuropsychological test performance in Obsessive-Compulsive disorder patients (n=21)


Resting state functional connectivity network		WCST	COWAT	SCWT	TMT-B	TMT-A	RCFT
<b>S3 vs. Rt. parahippocampal gyrus</b>	Correlation	0.137	-0.082	0.240	0.208	0.043	-0.069
	Significance	0.553	0.723	0.295	0.366	0.857	0.768
<b>S3 vs. Lt. associative visual cortex</b>	Correlation	-0.062	-0.068	-0.364	-0.187	0.065	0.064
	Significance	0.789	0.770	0.105	0.416	0.784	0.784
<b>S3 vs. Rt. Lingual gyrus</b>	Correlation	-0.144	0.316	-0.161	-0.302	-0.258	0.185
	Significance	0.533	0.163	0.487	0.183	0.273	0.422
<b>S5 vs. Rt. Fusiform gyrus</b>	Correlation	-0.154	0.375	-0.245	-0.211	-0.235	-0.266
	Significance	0.506	0.094	0.284	0.358	0.318	0.243
<b>S5 vs. Rt. Associative visual cortex</b>	Correlation	-0.358	0.207	-0.115	0.033	0.054	-0.252
	Significance	0.111	0.367	0.619	0.888	0.820	0.271
<b>S5 vs. Lt. primary somatosensory cortex</b>	Correlation	-0.368	-0.053	0.166	-0.092	-0.038	-0.108
	Significance	0.101	0.819	0.473	0.690	0.873	0.642

*Abbreviation:* Correlation= Pearson's correlation coefficient; COWAT= Category fluency score of the Controlled oral word association test; Lt.= left; RCFT= Organization score of the Rey-Osterrieth complex figure test; Rt.= right; TMT-A= Reaction time for Trail-making test A; TMT-B= Reaction time for the Trail-making test B; SCWT= Reaction time for the color-word page of Stroop color word test; WCST= Perseverative response score of Wisconsin card sorting test. \*p<.05 (two-tailed)

**Table 11.** Bivariate correlations between regional negative functional connectivity strength and neuropsychological test performance in healthy controls (n=22)

Resting state functional connectivity network		WCST	COWAT	SCWT	TMT-B	TMT-A	RCFT
<b>S3 vs. Rt. parahippocampal gyrus</b>	Correlation	-0.006	0.205	0.147	-0.253	-0.046	-0.282
	p value	0.979	0.360	0.515	0.255	0.837	0.204
<b>S3 vs. Lt. associative visual cortex</b>	Correlation	-0.184	0.141	0.126	-0.178	-0.194	-0.329
	p value	0.412	0.531	0.575	0.428	0.388	0.135
<b>S3 vs. Rt. Lingual gyrus</b>	Correlation	-0.228	0.578	0.059	-0.398	-0.03	-0.229
	p value	0.307	0.005*	0.794	0.066	0.895	0.305
<b>S5 vs. Rt. Fusiform gyrus</b>	Correlation	0.380	0.216	0.096	0.193	0.222	0.011
	p value	0.081	0.334	0.671	0.388	0.320	0.963
<b>S5 vs. Rt. Associative visual cortex</b>	Correlation	-0.139	0.258	-0.319	0	0.123	-0.126
	p value	0.538	0.247	0.147	>0.999	0.586	0.576
<b>S5 vs. Lt. primary somatosensory cortex</b>	Correlation	0.089	0.155	-0.158	0.195	-0.335	-0.188
	p value	0.694	0.490	0.481	0.385	0.127	0.402

*Abbreviation:* Correlation= Pearson's correlation coefficient; COWAT= Category fluency score of the Controlled oral word association test; Lt.= left; RCFT= Organization score of the Rey-Osterrieth complex figure test; Rt.= right; TMT-A= Reaction time for Trail-making test A; TMT-B= Reaction time for the Trail-making test B; SCWT= Reaction time for the color-word page of Stroop color word test; WCST= Perseverative response score of Wisconsin card sorting test. \*p<.05 (two-tailed)



**Figure 10.** Correlation between regional functional connectivity strength and neuropsychological test performance in obsessive-compulsive disorder. Stronger functional connectivity (a) between the ACC S5 seed region and the right claustrum predicted higher RCFT organization scores ( $r=0.541$ ,  $p$

= 0.011). And stronger functional connectivity (b) between ACC S7 seed region and the right supramarginal gyrus could predict shorter reaction time in TMT-A performance ( $r=-0.520$ ,  $p=0.019$ ). 95% confidence intervals are depicted with dotted lines.

*Abbreviations:* ACC= anterior cingulate cortex; RCFT= Rey-Osterreith Complex Figure Test; TMT-A= Trail Making Test, part A.

## DISCUSSION

This study demonstrated that each sub-region of the ACC had distinctive rs-FCNs in the OCD group, in which the strengths of the functional connectivity between ACC seeds versus specific components of the CSTC loop were altered. Our results also showed that altered rs-FC strength between the CSTC loop components and the ACC sub-regions was associated with executive dysfunction in OCD.

The CSTC loop model of OCD (71) proposes the involvement of direct and indirect cortico-striato-thalamic pathways, in which imbalances between these frontal–striatal circuits may mediate OCD symptomatology. The direct pathway is composed of the ventromedial frontal (ventral ACC and OFC)–ventral striatal (internal part of the globus pallidus) circuit, which results in thalamus disinhibition and thus an increased excitatory effect on the prefrontal cortex. In contrast, the indirect pathway is associated with a dorsal frontal (DLPFC and dorsal ACC)–dorsal striatal circuit and provides a mechanism of negative feedback, which is important for inhibition and switching of behavior. In this study, five seeds of the ACC, a central component of the CSTC loops in patients with OCD, constructed distinct patterns of rs-FCNs, in agreement with previous results in healthy subjects (69). However, between-group analyses revealed reduced rs-FCN strength in the claustrum, thalamus, and IPL in patients with OCD, which could be related to OCD symptomatology and impaired cognitive functioning.

Overall, each seeds of the ACC subregions constructed distinct patterns of rs-FCNs as a whole, consistent with the previous studies for healthy subjects which showed several distinct ACC-based rs-FCNs based on the functional segregations of ACC subregions, which are the motor control (S1), cognitive control (S3), conflict monitoring (S5), social processing (S7)



and emotional regulation (19) (69). In this study, the OCD group revealed reduced rs-FC strength in the brain areas which were indeed important components of the CSTC loop, namely the claustrum and the pulvinar of the thalamus (72), and in the inferior parietal lobule. As rs-FCNs reflect ongoing information processing between relevant brain areas during the task-free status and also shows regional brain activities during task performance(37), disruption of these rs-FCN strength and pattern could be a reflection of functional derangement in the relevant brain area.

We showed decreased regional functional connectivity between the ACC S1 seed and the left claustrum and between the ACC S5 seed and the right claustrum in the OCD group. The supragenual ACC, which includes the S5 seed, is typically associated with evaluative functions of reasoning (73), conflict detection (74, 75), and response to error (75). The claustrum, thin layers of gray matter tissue that reside between the insula and putamen, may receive, channel, and relay sensory information between several brain areas, namely many other neocortical areas in the frontal (76), temporal (77), and parietal cortices (78), as well as limbic structures such as the hippocampus (79), amygdala (80), caudate nucleus and putamen (81), comprising neural circuits subserving cognitive function (82) including the CSTC loops.

The RCFT task requires subjects initially to copy a complex geometric figure and then to reproduce that same figure based on the memory during the immediate and the delayed recall phase. As reproduction of that complex figure could be conducted only by the usage of organization strategies of frontal lobe functioning, the RCFT has been used as a valuable measure of executive functioning (12). In this study, we found no statistically significant differences in RCFT organization score between the patients with OCD and the healthy controls ( $p = 0.510$ ), which is in line with our previous studies (53, 83). However, stronger regional functional

connectivity between the S5 ACC seed and the right claustrum successfully predicted a higher RCFT organization score only in patients with OCD, raising the possibility that degrees of functional coherence between the supragenual ACC and the claustrum may be the neural correlate of the visual organization strategy in patients with OCD.

The subgenual ACC, which included Seed I9, is the central component of the limbic and paralimbic system, composed of the amygdala, insula, ventral striatum, and ventral regions of the ACC (84), subserves emotional responsiveness and regulation and the monitoring of rewarding or punishing outcomes (85). The subgenual ACC is involved in identifying the emotional significance of a stimulus and producing a relevant affective state as a response. The OFC is involved in the evaluation of motivational significance, generation of adaptive responses to rewarding or aversive stimuli, and regulation of emotional states. As a result, hyperactivity of OFC could disrupt the evaluative function for the consequences of immediate action, resulting in uncontrolled thoughts and behaviors (86). Considering these observations, an exceptionally strengthened rs-FC between the ACC I9 seed and OFC in the OCD group, which was revealed in this study, may be related to the strong anxiety precipitated by overvalued ideas regarding one's behaviors and thoughts.

This study showed that the weaker functional connectivity between the perigenual ACC S7 seed and the right IPL was significantly correlated with longer reaction times for completing the TMT-A task. Perigenual ACC, which includes the S7 ACC seed, has repeatedly been implicated in social cognitive functions, such as mentalizing (theory-of-mind) (87, 88) and self-reflection (89, 90). The IPL is a component of the so-called fronto-parietal control network, which consists of the DLPFC, ACC, and IPL, regulating cognitive control and decision-making processes (91). In this fronto-parietal control network, the IPL mainly supports proper task execution by taking

part in motor response preparation and visuospatial processing (92). Especially, the right IPL has been linked to the function of visuospatial memory, including the ability to sense relationships between body parts. It is also related to the perception of our own affects or feelings. In OCD patients, time-consuming repetitive compulsive acts and preoccupations to specific body parts or somatic discomforts could be related to the impaired self-reflection which underlies the functional impairment of the fronto-parietal control system. Some case reports have stressed the temporal relationships between OCD symptomatology and brain hemorrhage (93) and multiple sclerosis plaques (94) of the right parietal lobe. Similarly, disrupted structural connectivity (95) and altered chemical composition of white matter (96) in the supramarginal gyrus of the IPL subdivisions, have been linked to several executive dysfunctions including visuospatial ability and nonverbal memory as well as OCD symptom severity (97). Accordingly, defects in striato-fronto-parietal connectivity may underlie the inefficient information processing in OCD (98), which results in slowed psychomotor speed.

Relatively attenuated strength of rs-FC anti-correlation between ACC seeds and posterior brain areas should also be mentioned, as insufficient deactivation of specific brain areas in relation to ACC activation could result in inefficient information processing and impaired execution. Previous structural and functional neuroimaging studies revealed significantly lowered FA of lingual gyrus(95), excessive BOLD response of parahippocampal gyrus in decision of whether to discard personal items or not(99) and increased BOLD response in response to biological motion stimuli(100), which were in line with our study results.

There are some more issues to be discussed. This study did not consider OCD subtypes in the rs-FCN analyses. However, if we accept the concept of OCD as one disease entity with common pathophysiology encompassing

several symptom subtypes, rs-FC may reflect this common pathophysiology successfully. We also did not investigate rs-FCNs regarding left ACC seeds because previous studies of rs-FC for the ACC regions showed hemispheric similarity for the patterns of ACC connectivity (43, 101).

Our results, obtained from well-developed methodology, showed the importance of the ACC as a central executor in the CSTC loops of OCD (43). Using the rs-FCN approach, the patterns of which resemble the topographic activation patterns of the same regions during task performance (102), we provided evidence that the regional functional connectivity between ACC subregions and some components comprising the CSTC loops may be the neural correlates of executive dysfunction in OCD. Additionally, we revealed some beneficial clues pointing to the parietal cortex, particularly the IPL, as one of the important brain regions included in the OCD pathophysiology loops. Future research for causal relationships between brain regions related to the ACC rs-FC is needed.

## REFERENCES

1. Ruscio AM, Stein DJ, Chiu WT, Kessler RC. The epidemiology of obsessive-compulsive disorder in the National Comorbidity Survey Replication. *Mol Psychiatry*. 2010 Jan; 15(1):53-63.
2. Stewart SE, Rosario MC, Baer L, Carter AS, Brown TA, Scharf JM, et al. Four-factor structure of obsessive-compulsive disorder symptoms in children, adolescents, and adults. *J Am Acad Child Adolesc Psychiatry*. 2008 Jul; 47(7):763-72.
3. Abramowitz JS. The psychological treatment of obsessive-compulsive disorder. *Can J Psychiatry*. 2006 Jun; 51(7):407-16.
4. Gillan CM, Pappmeyer M, Morein-Zamir S, Sahakian BJ, Fineberg NA, Robbins TW, et al. Disruption in the balance between goal-directed behavior and habit learning in obsessive-compulsive disorder. *Am J Psychiatry*. 2011 Jul; 168(7):718-26.
5. Gottesman, II, Gould TD. The endophenotype concept in psychiatry: etymology and strategic intentions. *Am J Psychiatry*. 2003 Apr; 160(4):636-45.
6. Riesel A, Endrass T, Kaufmann C, Kathmann N. Overactive error-related brain activity as a candidate endophenotype for obsessive-compulsive disorder: evidence from unaffected first-degree relatives. *Am J Psychiatry*. 2011 Mar; 168(3):317-24.
7. Grisham JR, Anderson TM, Poulton R, Moffitt TE, Andrews G. Childhood neuropsychological deficits associated with adult obsessive-compulsive disorder. *Br J Psychiatry*. 2009 Aug; 195(2):138-41.
8. Viswanath B, Janardhan Reddy YC, Kumar KJ, Kandavel T, Chandrashekar CR. Cognitive endophenotypes in OCD: a study of unaffected siblings of probands with familial OCD. *Prog Neuropsychopharmacol Biol Psychiatry*. 2009 Jun 15; 33(4):610-5.

9. Cavedini P, Zorzi C, Piccinni M, Cavallini MC, Bellodi L. Executive dysfunctions in obsessive-compulsive patients and unaffected relatives: searching for a new intermediate phenotype. *Biol Psychiatry*. 2010 Jun 15; 67(12):1178-84.
10. Cummings JL. Anatomic and behavioral aspects of frontal-subcortical circuits. *Ann N Y Acad Sci*. 1995 Dec 15; 769:1-13.
11. Savage CR, Baer L, Keuthen NJ, Brown HD, Rauch SL, Jenike MA. Organizational strategies mediate nonverbal memory impairment in obsessive-compulsive disorder. *Biol Psychiatry*. 1999 Apr 1; 45(7):905-16.
12. Shin MS, Park SY, Park SR, Seol SH, Kwon JS. Clinical and empirical applications of the Rey-Osterrieth Complex Figure Test. *Nat Protoc*. 2006; 1(2):892-9.
13. Penades R, Catalan R, Andres S, Salamero M, Gasto C. Executive function and nonverbal memory in obsessive-compulsive disorder. *Psychiatry Res*. 2005 Jan 30; 133(1):81-90.
14. Bush G, Luu P, Posner MI. Cognitive and emotional influences in anterior cingulate cortex. *Trends Cogn Sci*. 2000 Jun; 4(6):215-22.
15. Vogt BA, Finch DM, Olson CR. Functional heterogeneity in c cingulate cortex: the anterior executive and posterior evaluative regions. *Cereb Cortex*. 1992 Nov-Dec; 2(6):435-43.
16. Gu BM, Park JY, Kang DH, Lee SJ, Yoo SY, Jo HJ, et al. Neural correlates of cognitive inflexibility during task-switching in obsessive-compulsive disorder. *Brain*. 2008 Jan; 131(Pt 1):155-64.
17. Ursu S, Stenger VA, Shear MK, Jones MR, Carter CS. Overactive action monitoring in obsessive-compulsive disorder: evidence from functional magnetic resonance imaging. *Psychol Sci*. 2003 Jul; 14(4):347-53.
18. Fitzgerald KD, Welsh RC, Gehring WJ, Abelson JL, Himle JA, Liberzon

- I, et al. Error-related hyperactivity of the anterior cingulate cortex in obsessive-compulsive disorder. *Biol Psychiatry*. 2005 Feb 1; 57(3):287-94.
19. Maltby N, Tolin DF, Worhunsky P, O'Keefe TM, Kiehl KA. Dysfunctional action monitoring hyperactivates frontal-striatal circuits in obsessive-compulsive disorder: an event-related fMRI study. *Neuroimage*. 2005 Jan 15; 24(2):495-503.
  20. Rotge JY, Guehl D, Dilharreguy B, Tignol J, Bioulac B, Allard M, et al. Meta-analysis of brain volume changes in obsessive-compulsive disorder. *Biol Psychiatry*. 2009 Jan 1; 65(1):75-83.
  21. Matsumoto R, Ito H, Takahashi H, Ando T, Fujimura Y, Nakayama K, et al. Reduced gray matter volume of dorsal cingulate cortex in patients with obsessive-compulsive disorder: a voxel-based morphometric study. *Psychiatry Clin Neurosci*. 2010 Oct; 64(5):541-7.
  22. Radua J, Mataix-Cols D. Voxel-wise meta-analysis of grey matter changes in obsessive-compulsive disorder. *Br J Psychiatry*. 2009 Nov; 195(5):393-402.
  23. Togao O, Yoshiura T, Nakao T, Nabeyama M, Sanematsu H, Nakagawa A, et al. Regional gray and white matter volume abnormalities in obsessive-compulsive disorder: a voxel-based morphometry study. *Psychiatry Res*. 2010 Oct 30; 184(1):29-37.
  24. Ha TH, Kang DH, Park JS, Jang JH, Jung WH, Choi JS, et al. White matter alterations in male patients with obsessive-compulsive disorder. *Neuroreport*. 2009 May 6; 20(7):735-9.
  25. Rotge JY, Guehl D, Dilharreguy B, Cuny E, Tignol J, Bioulac B, et al. Provocation of obsessive-compulsive symptoms: a quantitative voxel-based meta-analysis of functional neuroimaging studies. *J Psychiatry Neurosci*. 2008 Sep;33(5):405-12.
  26. Gu BM, Kang DH, Kwon JS. Understanding neuropsychiatric disorders:

- Insights from neuroimaging. New York: Cambridge university press; 2011. p. 247-59.
27. Ochsner KN, Bunge SA, Gross JJ, Gabrieli JD. Rethinking feelings: an fMRI study of the cognitive regulation of emotion. *J Cogn Neurosci*. 2002 Nov 15; 14(8):1215-29.
  28. Huyser C, Veltman DJ, Wolters LH, de Haan E, Boer F. Functional magnetic resonance imaging during planning before and after cognitive-behavioral therapy in pediatric obsessive-compulsive disorder. *J Am Acad Child Adolesc Psychiatry*. 2010 Dec;49(12): 1238-48, 48 e1-5.
  29. Cavanagh JF, Grundler TO, Frank MJ, Allen JJ. Altered cingulate sub-region activation accounts for task-related dissociation in ERN amplitude as a function of obsessive-compulsive symptoms. *Neuropsychologia*. 2010 Jun; 48(7):2098-109.
  30. Yucel M, Harrison BJ, Wood SJ, Fornito A, Wellard RM, Pujol J, et al. Functional and biochemical alterations of the medial frontal cortex in obsessive-compulsive disorder. *Arch Gen Psychiatry*. 2007 Aug; 64(8):946-55.
  31. Sumitani S, Harada M, Kubo H, Ohmori T. Proton magnetic resonance spectroscopy reveals an abnormality in the anterior cingulate of a subgroup of obsessive-compulsive disorder patients. *Psychiatry Res*. 2007 Jan 15; 154(1):85-92.
  32. Endrass T, Schuermann B, Kaufmann C, Spielberg R, Kniesche R, Kathmann N. Performance monitoring and error significance in patients with obsessive-compulsive disorder. *Biol Psychol*. 2010 May; 84(2):257-63.
  33. Carter CS, Braver TS, Barch DM, Botvinick MM, Noll D, Cohen JD. Anterior cingulate cortex, error detection, and the online monitoring of performance. *Science*. 1998 May 1; 280(5364):747-9.



34. Del Casale A, Kotzalidis GD, Rapinesi C, Serata D, Ambrosi E, Simonetti A, et al. Functional neuroimaging in obsessive-compulsive disorder. *Neuropsychobiology*. 2011; 64(2):61-85.
35. Friston K. Causal modelling and brain connectivity in functional magnetic resonance imaging. *PLoS Biol*. 2009 Feb 17; 7(2):e33.
36. Greicius MD, Supekar K, Menon V, Dougherty RF. Resting-state functional connectivity reflects structural connectivity in the default mode network. *Cereb Cortex*. 2009 Jan; 19(1):72-8.
37. Dosenbach NU, Fair DA, Cohen AL, Schlaggar BL, Petersen SE. A dual-networks architecture of top-down control. *Trends Cogn Sci*. 2008 Mar; 12(3):99-105.
38. Bar M. The proactive brain: using analogies and associations to generate predictions. *Trends Cogn Sci*. 2007 Jul; 11(7):280-9.
39. Ozaki TJ. Frontal-to-parietal top-down causal streams along the dorsal attention network exclusively mediate voluntary orienting of attention. *PLoS One*. 2011; 6(5):e20079.
40. Park CH, Chang WH, Ohn SH, Kim ST, Bang OY, Pascual-Leone A, et al. Longitudinal changes of resting-state functional connectivity during motor recovery after stroke. *Stroke*. 2011 May; 42(5):1357-62.
41. Cocchi L, Harrison BJ, Pujol J, Harding IH, Fornito A, Pantelis C, et al. Functional alterations of large-scale brain networks related to cognitive control in obsessive-compulsive disorder. *Hum Brain Mapp*. 2012 May; 33(5):1089-106.
42. Palomero-Gallagher N, Mohlberg H, Zilles K, Vogt B. Cytology and receptor architecture of human anterior cingulate cortex. *J Comp Neurol*. 2008 Jun 20; 508(6):906-26.
43. Margulies DS, Kelly AM, Uddin LQ, Biswal BB, Castellanos FX, Milham MP. Mapping the functional connectivity of anterior cingulate cortex. *Neuroimage*. 2007 Aug 15; 37(2):579-88.

44. Menzies L, Achard S, Chamberlain SR, Fineberg N, Chen CH, del Campo N, et al. Neurocognitive endophenotypes of obsessive-compulsive disorder. *Brain*. 2007 Dec; 130(Pt 12):3223-36.
45. Jang JH, Kim JH, Jung WH, Choi JS, Jung MH, Lee JM, et al. Functional connectivity in fronto-subcortical circuitry during the resting state in obsessive-compulsive disorder. *Neurosci Lett*. 2010 May 3; 474(3):158-62.
46. Meunier D, Ersche KD, Craig KJ, Fornito A, Merlo-Pich E, Fineberg NA, et al. Brain functional connectivity in stimulant drug dependence and obsessive-compulsive disorder. *Neuroimage*. 2012 Jan 16; 59(2):1461-8.
47. Fitzgerald KD, Welsh RC, Stern ER, Angstadt M, Hanna GL, Abelson JL, et al. Developmental alterations of frontal-striatal-thalamic connectivity in obsessive-compulsive disorder. *J Am Acad Child Adolesc Psychiatry*. 2011 Sep; 50(9):938-48 e3.
48. Goodman WK, Price LH, Rasmussen SA, Mazure C, Fleischmann RL, Hill CL, et al. The Yale-Brown Obsessive Compulsive Scale. I. Development, use, and reliability. *Arch Gen Psychiatry*. 1989 Nov; 46(11):1006-11.
49. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Archives of general psychiatry*. 1961 Jun; 4:561-71.
50. Beck AT, Steer R. Manual for the Beck Anxiety Inventory. San Antonio: Psychological Corporation; 1990.
51. Yum TH, Park YS, Oh KJ, Kim JG, Lee HY. The manual of Korean-Wechsler adult intelligence scale. Seoul: Korean Guidance Press; 1992.
52. Hashimoto N, Nakaaki S, Omori IM, Fujioi J, Noguchi Y, Murata Y, et al. Distinct neuropsychological profiles of three major symptom

- dimensions in obsessive-compulsive disorder. *Psychiatry Res.* 2011 May 15; 187(1-2):166-73.
53. Shin NY, Kang DH, Choi JS, Jung MH, Jang JH, Kwon JS. Do organizational strategies mediate nonverbal memory impairment in drug-naïve patients with obsessive-compulsive disorder? *Neuropsychology.* 2010 Jul; 24(4):527-33.
54. Eling P, Derckx K, Maes R. On the historical and conceptual background of the Wisconsin Card Sorting Test. *Brain Cogn.* 2008 Aug; 67(3):247-53.
55. Heaton RK, Chelune GJ, Talley JL, Kay GG, Curtiss G. *Wisconsin Card Sorting Test manual: revised and expanded.* Odessa, Florida: Psychological Assessment Resources; 1993.
56. Greve KW. Can perseverative responses on the Wisconsin Card Sorting Test be scored accurately? *Arch Clin Neuropsychol.* 1993 Nov; 8(6):511-7.
57. Tükel R, Gurvit H, Ertekin BA, Oflaz S, Ertekin E, Baran B, et al. Neuropsychological function in obsessive-compulsive disorder. *Compr Psychiatry.* 2012 Feb; 53(2):167-75.
58. Sumner JA, Griffith JW, Mineka S. Examining the mechanisms of overgeneral autobiographical memory: capture and rumination, and impaired executive control. *Memory.* 2011 Feb; 19(2):169-83.
59. Golden CJ, Freshwater SM. *The stroop color and word test: A manual for clinical and experimental uses.* Chicago, IL: Stoelting Co; 2002.
60. Reitan RM. Validity of the Trail Making Test as an indicator of organic brain damage. *Perceptual and Motor Skills.* 1958; 8:271-6.
61. Rey A. L'examen psychologique dans les cas d'encephalopathie traumatique. *Arch Psychol.* 1941; 28:286-340.
62. Osterrieth PA. Le test de copie d'une figure complexe. *Arch Psychol.* 1944; 30:206-356.

63. Somerville J, Tremont G, Stern RA. The Boston Qualitative Scoring System as a measure of executive functioning in Rey-Osterrieth Complex Figure performance. *J Clin Exp Neuropsychol*. 2000 Oct; 22(5):613-21.
64. Bennett-Levy J. Determinants of performance on the Rey-Osterrieth Complex Figure Test: an analysis, and a new technique for single-case assessment. *Br J Clin Psychol*. 1984 May; 23 (Pt 2):109-19.
65. Stern RA, Javorsky DJ, Singer EA, Singer NG, Somerville JA, Duke LM. The Boston Qualitative Scoring System for the Rey-Osterrieth Complex Figure. Odessa: Psychological Assessment Resources; 1999.
66. Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME. The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proc Natl Acad Sci U S A*. 2005 Jul 5; 102(27):9673-8.
67. Birn RM, Diamond JB, Smith MA, Bandettini PA. Separating respiratory-variation-related fluctuations from neuronal-activity-related fluctuations in fMRI. *Neuroimage*. 2006 Jul 15; 31(4):1536-48.
68. Macey PM, Macey KE, Kumar R, Harper RM. A method for removal of global effects from fMRI time series. *Neuroimage*. 2004 May; 22(1):360-6.
69. Kelly AM, Di Martino A, Uddin LQ, Shehzad Z, Gee DG, Reiss PT, et al. Development of anterior cingulate functional connectivity from late childhood to early adulthood. *Cereb Cortex*. 2009 Mar; 19(3):640-57.
70. Bush G, Vogt BA, Holmes J, Dale AM, Greve D, Jenike MA, et al. Dorsal anterior cingulate cortex: a role in reward-based decision making. *Proc Natl Acad Sci U S A*. 2002 Jan 8; 99(1):523-8.
71. Saxena S, Bota RG, Brody AL. Brain-behavior relationships in obsessive-compulsive disorder. *Semin Clin Neuropsychiatry*. 2001 Apr; 6(2):82-101.

72. Kwon JS, Jang JH, Choi JS, Kang DH. Neuroimaging in obsessive-compulsive disorder. *Expert Rev Neurother*. 2009 Feb; 9(2):255-69.
73. Luo Q, Perry C, Peng D, Jin Z, Xu D, Ding G, et al. The neural substrate of analogical reasoning: an fMRI study. *Brain Res Cogn Brain Res*. 2003 Oct; 17(3):527-34.
74. Botvinick MM, Cohen JD, Carter CS. Conflict monitoring and anterior cingulate cortex: an update. *Trends Cogn Sci*. 2004 Dec; 8(12):539-46.
75. Lutcke H, Frahm J. Lateralized anterior cingulate function during error processing and conflict monitoring as revealed by high-resolution fMRI. *Cereb Cortex*. 2008 Mar; 18(3):508-15.
76. Kievit J, Kuypers HG. Subcortical afferents to the frontal lobe in the rhesus monkey studied by means of retrograde horseradish peroxidase transport. *Brain Res*. 1975 Feb 28; 85(2):261-6.
77. Webster MJ, Bachevalier J, Ungerleider LG. Transient subcortical connections of inferior temporal areas TE and TEO in infant macaque monkeys. *J Comp Neurol*. 1995 Feb 6; 352(2):213-26.
78. Pearson RC, Brodal P, Gatter KC, Powell TP. The organization of the connections between the cortex and the claustrum in the monkey. *Brain Res*. 1982 Feb 25; 234(2):435-41.
79. Amaral DG, Cowan WM. Subcortical afferents to the hippocampal formation in the monkey. *J Comp Neurol*. 1980 Feb 15; 189(4):573-91.
80. Amaral DG, Insausti R. Retrograde transport of D-[3H]-aspartate injected into the monkey amygdaloid complex. *Exp Brain Res*. 1992; 88(2):375-88.
81. Arikuni T, Kubota K. Claustral and amygdaloid afferents to the head of the caudate nucleus in macaque monkeys. *Neurosci Res*. 1985 Apr; 2(4):239-54.
82. Edelstein LR, Denaro FJ. The claustrum: a historical review of its anatomy, physiology, cytochemistry and functional significance. *Cell*

- Mol Biol (Noisy-le-grand). 2004 Sep; 50(6):675-702.
83. Park HS, Shin YW, Ha TH, Shin MS, Kim YY, Lee YH, et al. Effect of cognitive training focusing on organizational strategies in patients with obsessive-compulsive disorder. *Psychiatry Clin Neurosci*. 2006 Dec; 60(6):718-26.
  84. Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion perception I: The neural basis of normal emotion perception. *Biol Psychiatry*. 2003 Sep 1; 54(5):504-14.
  85. Knutson B, Fong GW, Bennett SM, Adams CM, Hommer D. A region of mesial prefrontal cortex tracks monetarily rewarding outcomes: characterization with rapid event-related fMRI. *Neuroimage*. 2003 Feb; 18(2):263-72.
  86. Aouizerate B, Guehl D, Cuny E, Rougier A, Bioulac B, Tignol J, et al. Pathophysiology of obsessive-compulsive disorder: a necessary link between phenomenology, neuropsychology, imagery and physiology. *Prog Neurobiol*. 2004 Feb; 72(3):195-221.
  87. Frith U, Frith CD. Development and neurophysiology of mentalizing. *Philos Trans R Soc Lond B Biol Sci*. 2003 Mar 29; 358(1431):459-73.
  88. Amodio DM, Frith CD. Meeting of minds: the medial frontal cortex and social cognition. *Nat Rev Neurosci*. 2006 Apr; 7(4):268-77.
  89. Ochsner KN, Beer JS, Robertson ER, Cooper JC, Gabrieli JD, Kihlstrom JF, et al. The neural correlates of direct and reflected self-knowledge. *Neuroimage*. 2005 Dec; 28(4):797-814.
  90. Johnson SC, Baxter LC, Wilder LS, Pipe JG, Heiserman JE, Prigatano GP. Neural correlates of self-reflection. *Brain*. 2002 Aug; 125(Pt 8):1808-14.
  91. Vincent JL, Kahn I, Snyder AZ, Raichle ME, Buckner RL. Evidence for a frontoparietal control system revealed by intrinsic functional connectivity. *J Neurophysiol*. 2008 Dec; 100(6):3328-42.

92. Lazeron RH, Rombouts SA, Machielsen WC, Scheltens P, Witter MP, Uylings HB, et al. Visualizing brain activation during planning: the tower of London test adapted for functional MR imaging. *AJNR Am J Neuroradiol*. 2000 Sep; 21(8):1407-14.
93. Ros LT, Podgorski JK. Obsessive-compulsive disorder secondary to a right parietal lobe haemorrhage: a case report. *Acta Neurol Belg*. 2007 Sep; 107(3):94-5.
94. Douzenis A, Michalopoulou PG, Voumvourakis C, Typaldou M, Michopoulos I, Lykouras L. Obsessive-compulsive disorder associated with parietal white matter multiple sclerosis plaques. *World J Biol Psychiatry*. 2009; 10(4 Pt 3):956-60.
95. Szeszko PR, Ardekani BA, Ashtari M, Malhotra AK, Robinson DG, Bilder RM, et al. White matter abnormalities in obsessive-compulsive disorder: a diffusion tensor imaging study. *Arch Gen Psychiatry*. 2005 Jul; 62(7):782-90.
96. Menzies L, Williams GB, Chamberlain SR, Ooi C, Fineberg N, Suckling J, et al. White matter abnormalities in patients with obsessive-compulsive disorder and their first-degree relatives. *Am J Psychiatry*. 2008 Oct; 165(10):1308-15.
97. Kwon JS, Kim JJ, Lee DW, Lee JS, Lee DS, Kim MS, et al. Neural correlates of clinical symptoms and cognitive dysfunctions in obsessive-compulsive disorder. *Psychiatry Res*. 2003 Jan 20; 122(1):37-47.
98. Okasha A, Rafaat M, Mahallawy N, El Nahas G, El Dawla AS, Sayed M, et al. Cognitive dysfunction in obsessive-compulsive disorder. *Acta Psychiatr Scand*. 2000 Apr; 101(4):281-5.
99. Tolin DF, Kiehl KA, Worhunsky P, Book GA, Maltby N. An exploratory study of the neural mechanisms of decision making in compulsive hoarding. *Psychol Med*. 2009 Feb; 39(2):325-36.

100. Jung WH, Gu BM, Kang DH, Park JY, Yoo SY, Choi CH, et al. BOLD response during visual perception of biological motion in obsessive-compulsive disorder : an fMRI study using the dynamic point-light animation paradigm. *Eur Arch Psychiatry Clin Neurosci*. 2009 Feb; 259(1):46-54.
101. Yu C, Zhou Y, Liu Y, Jiang T, Dong H, Zhang Y, et al. Functional segregation of the human cingulate cortex is confirmed by functional connectivity based neuroanatomical parcellation. *Neuroimage*. 2011 Feb 14; 54(4):2571-81.
102. Deco G, Corbetta M. The dynamical balance of the brain at rest. *Neuroscientist*. 2011 Feb; 17(1):107-23.



## 국문 초록

**서론:** 앞쪽 띠피질은 강박증의 유력한 신경해부학적 가설인 피질-피질하 신경회로 이상 모델을 구성하는 중요한 뇌부위로서, 강박증 환자에서 앞쪽 띠피질의 기능 이상은 강박 증상 및 실행기능의 손상과도 관련되어 있을 가능성이 제기되어 왔다. 본 연구에서는 운동기능 조절, 주의력 통제, 상반된 정보들을 바탕으로 한 의사결정 및 감정조절 등 각각의 수행 기능에 따라 앞쪽 띠피질을 다섯 개의 세부 영역으로 나눈 후, 이러한 세부 영역 각 부위들과 다른 뇌 부위들 사이의 휴지기 기능적 뇌연결성을 관찰하고 정상인의 경우와 비교한 후, 강박증 환자에서 앞쪽 띠피질과 다른 뇌부위들 사이의 휴지기 기능적 뇌연결성의 변화가 다양한 실행기능의 손상 사이의 관계를 규명하고자 하였다.

**방법:** 본 연구에서는 이전에 정신과 약물 투약력이 없거나 혹은 연구 참여 당시로부터 최근 4주 이상의 기간 동안 정신과 약물을 투약하지 않은 21명의 강박증 환자와 22명의 정상 대조군을 대상으로 하여, 특정한 과제를 수행하지 않는 휴식 상태에서 기능적 뇌 자기공명영상을 촬영하였다. 이와 같이 얻어진 영상을 이용하여, 앞쪽 띠피질의 전장에 걸쳐 균일한 간격으로 분포된 관심 영역 중, 이전의 기능적 뇌영상 연구를 통해 앞쪽 띠피질이 수행하는 다양한 실행기능과의 관련성이 시사되는 다섯 개의 관심 영역을 선정하고, 이들 관심 영역들 각각과 이외의 다른 뇌 영역들 사이에 구성되는 휴지기 기능적 뇌연결성의 양상을 탐색하였다. 더불어, 강박증 환자군과 정상 대조군 사이에서 휴지기 기능적 뇌연결성의 강도가 유의한 차이를 보이는 경우, 앞쪽

띠피질의 관심영역과 해당 뇌영역 사이 기능적 연결성의 강도와 실행기능 손상 정도 사이의 관련성을 검증하였다.

**결과:** 강박증 환자와 정상 대조군 모두에서, 앞쪽 띠피질의 문층으로부터 미층까지 분포하는 다섯 개의 관심영역들은 각기 서로 다른 형태의 휴지기 기능적 뇌연결성을 보이는 공통점이 있었다. 두 군을 비교하였을 때, 정상 대조군에 비해 강박증 환자군은 앞쪽 띠피질과 시상, 하측 두정엽 및 담장(claustrum)과의 휴지기 기능적 뇌연결성이 유의하게 감소되어 있었다. 이 중 앞쪽 띠피질의 무릎 위쪽 부위에 위치한 S5 관심영역과 담장 사이의 기능적 연결성이 강할수록 레이 복합도형검사의 구조화 점수가 우수하였고, 앞쪽 띠피질의 무릎 밑쪽 부위에 위치한 S7 관심영역과 우반구의 하측 두정엽 사이의 기능적 연결성이 강할수록 길 만들기 검사 A형 수행시의 수행시간이 짧아지는 양상이 관찰되었다.

**결론:** 본 연구의 결과는 강박증의 피질-피질하 신경모델을 구성하는 앞쪽 띠피질과 다른 뇌 영역들 사이의 휴지기 기능적 뇌연결성 변화가 강박증에서의 실행기능 손상과 관련되어 있음을 보여 준다. 또한 전전두엽의 영역들뿐만 아니라 하측 두정엽 또한 강박증의 피질-피질하 신경모델에서 정보처리 속도 등과 관련하여 중요한 구성 요소로 작용할 가능성을 시사한다.

**주요어:** 앞쪽 띠피질, 휴지기 기능적 뇌연결성, 기능적 뇌 자기공명영상, 강박증, 집행기능

**학번:** 2010-23694